



Indeno[1,2,3-cd]pyrene

Environmental estimates (circa 2011): Supplemental data

Table of Contents

1. Data for lifetime excess cancer risk estimates.....	2
Overview	2
i. Environmental Concentrations.....	2
ii. Calculated Lifetime Daily Intake	2
iii. Cancer Potency Factors	2
iv. Lifetime Excess Cancer Risk (per million people)	3
Supporting data by exposure pathway	3
i. Outdoor air	3
ii. Indoor air	3
iii. Dust.....	5
iv. Drinking water	6
v. Food and Beverages	6
2. Data quality for lifetime excess cancer risk estimates	9
3. Data for mapping concentrations	10
Estimates by health region	10
Estimates by census block	10

1. Data for lifetime excess cancer risk estimates

Overview

The summary data used to calculate lifetime excess cancer risk and the results for indeno(1,2,3-cd)pyrene are provided in the tables below. For more detailed information on supporting data and sources, see below for each exposure pathway.

i. Environmental Concentrations

Exposure pathway	Units	Average	Maximum	Notes
Outdoor air	µg/m ³	0.0001	0.00058	
Indoor air	µg/m ³	0.00044	0.0018	
Dust	µg/g	3.07	33.5	
Drinking water	µg/L	Insufficient data		
Foods and beverages		See detailed data	Not estimated	

ii. Calculated Lifetime Daily Intake

Exposure pathway	Average intake (mg/kg bodyweight per day)	Maximum intake (mg/kg bodyweight per day)
Outdoor air	0.000000023	0.00000013
Indoor air	0.00000014	0.00000058
Dust	0.0000020	0.000022
Drinking water	Insufficient data	
Foods and beverages	0.0000001	Not estimated

iii. Cancer Potency Factors

Exposure route	Health Canada	US EPA	CA OEHHA
Inhalation	--	--	0.39
Ingestion	--	--	1.2

Sources for Cancer Potency Factors:

- Health Canada, 2010. Federal Contaminated Site Risk Assessment in Canada, Part I: Guidance on Human Health Preliminary Quantitative Risk Assessment. Version 2.0.
- Health Canada, 2010. Federal Contaminated Site Risk Assessment in Canada, Part II: Health Canada Toxicological Reference Values (TRVs) and Chemical-Specific Factors. Version 2.0.
- United States Environmental Protection Agency Integrated Risk Information System
- California Office of Environmental Health Hazard Assessment, 2009. Air Toxics Hot Spots Risk Assessment Guidelines Part II: Technical Support Document for Cancer Potency Factors, Appendix A. (Updated 2011)

iv. Lifetime Excess Cancer Risk (per million people)

Exposure pathway	Average ¹			Maximum ²
	Health Canada	US EPA	CA OEHHA ³	
Outdoor air	--	--	0.0009	0.0053
Indoor air	--	--	0.056	0.23
Dust	--	--	2.42	26.40
Drinking water	Insufficient data			
Foods and beverages	--	--	--	Not estimated

¹Lifetime excess cancer risk based on average intake x cancer potency factor from each agency

²Lifetime excess cancer risk based on maximum intake x highest cancer potency factor

³California Office of Environmental Health Hazard Assessment

Supporting data by exposure pathway

i. Outdoor air

Outdoor air concentrations are from the National Air Pollution Surveillance monitoring network operated by Environment Canada, for the year 2010.

Source	Stations (n)	Min	Max	Mean	DF
NAPS 2010 ($\mu\text{g}/\text{m}^3$)	17	0.000019	0.00058	0.0001	1.0

DF = Detection frequency

We assume indeno(1,2,3-cd)pyrene is present at these levels in all outdoor air, although concentrations may vary from one location to another.

ii. Indoor air

Indoor air concentrations are based on data published in peer-reviewed literature since 2000. A ranking system was used to select data most representative of Canadian conditions circa 2011:

1. Canadian data collected in 2000 or more recently, sample duration of 24 hours or longer;
2. US studies of similar currency and sample duration;
3. Studies from northern European countries of similar currency and sample duration;
4. Canadian, US or European studies with data collected prior to 2000 and similar sample duration; and
5. Studies with sample duration of less than 24 hours regardless of country or collection date, or studies from countries not comparable to Canada.

Rank:	2	Author:	Jung (2010)	Location:	New York City						
Samples (n)	DF*	DL**	Sample Date	Units	Sample Duration	Min	Max	Mean (AM)	Med	Geomean (GM)	Percentile
203			2005 -	µg/m ³	14 days	0.00005	0.0022	0.000435	0.00035	0.00026	
98			2010			0.00002	0.001	0.0004	0.00016		

Notes: Values listed in the following order: heating season (Oct-Apr), non-heating season (May-Sept)
 *DF = Detection frequency
 **DL = Detection limit

Rank:	2	Author:	Li (2005)	Location:	Chicago						
Samples (n)	DF*	DL**	Sample Date	Units	Sample Duration	Min	Max	Mean (AM)	Med	Geomean (GM)	Percentile
10	~0.95		2000 - 2001	µg/m ³	48h x 14 months	0.000001	0.001		0.00045		10th 0.000001 25th 0.00002 75th 0.0003 90th 0.0005

Notes: non-smoking homes, (sampled once a month for 14 months) total n = 115
 *DF = Detection frequency
 **DL = Detection limit

Rank:	3	Author:	Gustafson (2008)	Location:	Hagorfs, Sweden						
Samples (n)	DF*	DL**	Sample Date	Units	Sample Duration	Min	Max	Mean (AM)	Med	Geomean (GM)	Percentile
13	1	0.00005	Feb-	µg/m ³	24hr	0.00005	0.0028	0.00061	0.00028		
10	0.8		March 2003			<DL	0.00017	0.00010	0.00010		

Notes: Values listed in the following order: wood-burning, non-wood burning homes.
 *DF = Detection frequency
 **DL = Detection limit

Rank:	4	Author:	Sanderson (2004)	Location:	Beauharnois, Quebec						
Samples (n)	DF*	DL**	Sample Date	Units	Sample Duration	Min	Max	Mean (AM)	Med	Geomean (GM)	Percentile
3	0.84	0.00001		µg/m ³	24h			0.00031			
12								0.0002		0.000073	

Notes: near aluminum smelter, Values listed in the following order: homes with oil heating, with no oil heating
 *DF = Detection frequency
 **DL = Detection limit

Sources for indoor air data:

- Gustafson P, Östman C, Sällsten G. 2008. Indoor levels of polycyclic aromatic hydrocarbons in homes with or without wood burning for heating. *Environ Sci Technol* 42: 5074-5080.
- Jung K, Patel MM, Kinney PL, Chillrud SN, Whyatt R, Hoepner L, et al. 2010. 1. Effects of Season and Indoor Heating on Indoor and Outdoor Residential Levels of Airborne Polycyclic Aromatic Hydrocarbons, Absorbance and Particulate Matter 2.5 in an Inner City Cohort of Young Children. *Journal of Allergy and Clinical Immunology* 125: AB81.

- Li A, Schoonover TM, Zou QM, Norlock F, Conroy LM, Scheff PA, et al. 2005. Polycyclic aromatic hydrocarbons in residential air of ten Chicago area homes: Concentrations and influencing factors. Atmospheric Environment 39: 3491-3501.
- Sanderson EG, Farant JP. 2004. Indoor and outdoor polycyclic aromatic hydrocarbons in residences surrounding a Soderberg aluminum smelter in Canada. Environ Sci Technol 38: 5350-5356.

iii. Dust

Indoor dust concentrations are based on data published in peer-reviewed literature since 2000. A ranking system was used to select data most representative of Canadian conditions circa 2011:

1. Canadian data collected in 2000 or more recently, sample duration of 24 hours or longer;
2. US studies of similar currency and sample duration;
3. Studies from northern European countries of similar currency and sample duration;
4. Canadian, US or European studies with data collected prior to 2000 and similar sample duration; and
5. Studies with sample duration of less than 24 hours regardless of country or collection date, or studies from countries not comparable to Canada.

Rank:	1	Author:	Maertens (2008)				Location:	Ottawa, Canada				
Samples (n)	DF*	DL**	Sample Date	Units	Sample Duration	Min	Max	Mean (AM)	Med	Geomean (GM)	Percentile	
51	1.0	0.039	2002 - 2003	µg/g		0.1	33.5	3.07	0.91	1.29		

Notes: Analyzed using GC/MS

*DF = Detection frequency

**DL = Detection limit

Rank:	2	Author:	Whitehead (2011)				Location:	California, USA				
Samples (n)	DF*	DL**	Sample Date	Units	Sample Duration	Min	Max	Mean (AM)	Med	Geomean (GM)	Percentile	
583	0.998	0.002	2001-2007	µg/g		<DL	2.371		0.053			

Notes: Analyzed using GC/MS

*DF = Detection frequency

**DL = Detection limit

Rank:	2	Author:	Hoh (2012)				Location:	San Diego County, CA, USA				
Samples (n)	DF*	DL**	Sample Date	Units	Sample Duration	Min	Max	Mean (AM)	Med	Geomean (GM)	Percentile	
43	1.0		2005-2007	µg/g		0.00482	0.171		0.0365		25 th 0.0224 75 th 0.0593	
89	1.0					<DL	0.528		0.0586		25 th 0.0310 75 th 0.0949	

Notes: Analyzed using GC/MS

*DF = Detection frequency

**DL = Detection limit

Sources for dust:

- Hoh E, Hunt RN, Quintana PJE, Zakarian JM, Chatfield DA, Wittry BC, Rodriguez E, Matt GE. 2012. Environmental tobacco smoke as a source of polycyclic aromatic hydrocarbons in settled house dust. *Environ Sci Technol* 46: 4174-4183.
- Maertens RM, Yang XF, Zhu JP, Gagne RW, Douglas GR, White PA. 2008. Mutagenic and carcinogenic hazards of settled house dust I: Polycyclic aromatic hydrocarbon content and excess lifetime cancer risk from preschool exposure. *Environmental Science & Technology* 42: 1747-1753.
- Whitehead T, Metayer C, Gunier RB, Ward MH, Nishioka MG, Buffler P, Rappaport SM. 2011. Determinants of polycyclic aromatic hydrocarbon levels in house dust. *J Expo Sci Environ Epidemiol* 21(2): 123-132.

iv. Drinking water

No recent data or studies were identified.

v. Food and Beverages

Food consumption data are from the Statistics Canada Food Statistics Report (2006) - Food available, adjusted for losses tables, and from the Nutrition Canada Survey (1970-1972).

Food concentration data are from the US Total Diet Study (2003-2004), and the US EPA's Dietary Exposure Potential Models (v5.0 2003), with the exception of data on metals, which are from the Canadian Food Inspection Agency (2004-2005).

In order to better represent actual intake, we incorporated data for cooked and/or processed foods, as in some cases this can either add to or diminish the amounts of a substance measured in raw food.

Concentration data were obtained for 54% of total seafood consumed.

Food or Beverage	Concentration (µg/g)	DF	Food or Beverage	Concentration (µg/g)	DF
Beef			Peaches fresh		
Chicken			Pears canned		
Mutton and lamb			Pears fresh		
Offal			Pineapples canned		
Oils and fats			Pineapples fresh		
Pork			Plums total fresh		
Salad oils			Quinces fresh		
Shortening and shortening oils			Raspberries frozen		
Stewing hen			Strawberries canned		
Turkey			Strawberries fresh		
Veal			Strawberries frozen		
Fish fresh and frozen seafood	0.04671	0.0157	Sugar maple		
Fish freshwater	0.04168	0.0132	Sugar refined		
Fish processed seafood			Honey		
Apple pie filling			Artichokes fresh		
Apple sauce			Asparagus canned		
Apples canned			Asparagus fresh		
Apples dried			Avocados fresh		
Apples fresh			Beans baked and canned		
Apples frozen			Beans dry		
Apricots canned			Beans green and wax canned		
Apricots fresh			Beans green and wax fresh		
Bananas fresh			Beans green and wax frozen		
Berries other fresh			Beets canned		
Blueberries canned			Beets fresh		
Blueberries fresh			Broccoli fresh		
Blueberries frozen			Broccoli frozen		
Cherries fresh			Brussels sprouts fresh		
Cherries frozen			Brussels sprouts frozen		
Citrus other fresh			Cabbage Chinese fresh		
Coconut fresh			Cabbage fresh		
Cranberries fresh			Carrots canned		
Dates fresh			Carrots fresh		
Figs fresh			Carrots frozen		
Fruit dried			Cauliflower fresh		
Grapefruit fresh			Cauliflower frozen		
Grapes fresh			Celery fresh		
Guava and mangoes fresh			Corn canned		
Kiwi fresh			Corn flour and meal		
Lemons fresh			Corn fresh		
Limes fresh			Corn frozen		
Mandarins fresh			Cucumbers fresh		
Melons musk, cantaloupe fresh			Eggplant fresh		
Melons other fresh			Garlic fresh		
Melons watermelons fresh			Kohlrabi fresh		
Melons, winter melons fresh			Leeks fresh		
Nectarines fresh			Lettuce fresh		
Oranges fresh			Lima beans frozen		
Papayas fresh			Manioc fresh		
Peaches canned			Mushrooms canned		

Food or Beverage	Concentration (µg/g)	DF
Mushrooms fresh		
Okra fresh		
Olives fresh		
Onions and shallots fresh		
Parsley fresh		
Parsnips fresh		
Peas canned		
Peas dry		
Peas fresh		
Peas frozen		
Peppers fresh		
Potatoes chips		
Potatoes frozen		
Potatoes other processed		
Potatoes sweet fresh		
Potatoes white fresh		
Potatoes white fresh and processed		
Pumpkins and squash fresh		
Radishes fresh		
Rappini fresh		
Rutabagas and turnip fresh		
Spinach fresh		
Spinach frozen		
Tomatoes canned		
Tomatoes fresh		
Tomatoes pulp, paste and puree		
Vegetables other edible root fresh		
Vegetables other leguminous fresh		
Vegetables unspecified canned		
Vegetables unspecified fresh		
Vegetables unspecified frozen		
Butter		
Cheese cheddar		
Cheese cottage		
Cheese processed		
Cheese variety		
Cream cereal 10%		
Cream sour		
Cream table 18%		
Cream whipping 32% or 35%		
Eggs		
Ice cream		
Ice milk		
Margarine		

Food or Beverage	Concentration (µg/g)	DF
Milk buttermilk		
Milk chocolate drink		
Milk concentrated skim		
Milk concentrated whole		
Milk other whole milk products		
Milk partly skimmed 2%		
Milk skim		
Milk standard		
Milk sweetened concentrated skim		
Milkshake		
Powder buttermilk		
Powder skim milk		
Powder whey		
Sherbet		
Yogurt		
Cereal products		
Oatmeal and rolled oats		
Peanuts		
Pot and pearl barley		
Pulses and nuts		
Rice		
Rye flour		
Tree nuts		
Wheat flour		
Ale, beer, stout and porter		
Beverages alcoholic		
Coffee		
Distilled spirits		
Juice apple		
Juice grape		
Juice tomato		
Juice fruit		
Juice grapefruit		
Juice lemon		
Juice orange		
Juice pineapple		
Juice vegetable		
Soft drinks		
Tea		
Water bottled		
Wines		
Cocoa		

2. Data quality for lifetime excess cancer risk estimates

Only publicly available data were used to calculate these indicators. Data that are not publicly available may produce different results.

No systematic method for measuring data quality was possible, so we provide the following assessments of how well the data used may represent the actual Canadian average levels. Quality is rated higher when there are data from a number of Canadian monitors, or from Canadian studies that show results similar to other comparable studies. Quality is rated lower when data from few monitors or studies were available, and lowest when estimates are based on non-Canadian data. Others may rate data quality differently.

Exposure Pathway	Data Quality	Notes
Outdoor air	Low	<ul style="list-style-type: none"> Indeno(1,2,3-cd)pyrene is regularly measured in outdoor air at 17 monitoring stations across Canada using accepted protocols.
Indoor air	Very Low	<ul style="list-style-type: none"> One recent US study identified (New York City). Agrees reasonably well with a smaller US study in Chicago. A small Canadian study near an aluminum smelter reported a lower mean for 10 homes without oil heating, but a similar mean for 3 homes with oil heating.
Indoor dust	Low	<ul style="list-style-type: none"> Measured levels from one recent Canadian study (Ottawa, ON) are considerably higher than 2 recent studies conducted in California, USA using the same analytical methods.
Drinking water	Gap	<ul style="list-style-type: none"> Only 1 sample was analyzed for indeno(1,2,3-cd)pyrene in Ontario in 2009. No recent data or studies were identified.
Foods and beverages	Gap	<ul style="list-style-type: none"> No Canadian or US data on concentrations of indeno(1,2,3-cd)pyrene in foods or beverages were identified.

3. Data for mapping concentrations

The maps use geographic coordinates at the census block level to represent residential locations. Concentration estimates are mapped at the health region level, which are created with aggregated census block data.

We used a model to predict annual average concentrations of indeno(1,2,3-cd)pyrene in outdoor air at residential locations for 2011. These are predicted using levels measured from the National Air Pollution Surveillance (NAPS) monitors and estimated concentrations from known emitters. For more information on how these estimates were created, please see the Mapping Methods document on the [Environmental Approach](#) section of our website.

Estimates by health region

The table below shows predicted indeno(1,2,3-cd)pyrene concentrations by province based on data at the health region level. The median concentration of indeno(1,2,3-cd)pyrene measured in outdoor air in 2011 at the health region level was 0.00023 $\mu\text{g}/\text{m}^3$, while the mean concentration was 0.00026 $\mu\text{g}/\text{m}^3$. Concentrations of indeno(1,2,3-cd)pyrene can be higher or lower than average in many locations.

i. Provincial averages of predicted indeno(1,2,3-cd)pyrene concentrations ($\mu\text{g}/\text{m}^3$) in outdoor air in 2011 based on health regions

Province	Median	Mean
BC	0.00029	0.00029
AB	0.00017	0.00017
SK	0.00015	0.00016
MB	0.00015	0.00015
ON	0.00025	0.00031
QC	0.00025	0.00032
NB	0.00023	0.00022
PE	0.00022	0.00022
NS	0.00027	0.00027
NL	0.00014	0.00015
YK	0.00021	0.00021
NT	0.00015	0.00015
NU	0.00025	0.00025
Canada	0.00023	0.00026

Estimates by census block

The table below shows provincial populations by concentration levels (either annual average or number of times above/below the national average) based on the census block data and the associated potential lifetime excess risk given different cancer potency factors.

i. Provincial population distribution by estimated average concentration ($\mu\text{g}/\text{m}^3$) of indeno(1,2,3-cd)pyrene in outdoor air in 2011 based on NAPS data at the census block

Estimated annual average concentration ($\mu\text{g}/\text{m}^3$)	Less than 0.000033	0.000033 to 0.00004	0.00004 to 0.00005	0.00005 to 0.000067	0.000067 to 0.0001	0.0001 to 0.00015	0.00015 to 0.0002	0.0002 to 0.00025	0.00025 to 0.0003	More than 0.0003
	> 3x lower	2.5 to 3x lower	2 to 2.5x lower	1.5 to 2x lower	1 to 1.5x lower	1 to 1.5x higher	1.5 to 2x higher	2 to 2.5x higher	2.5 to 3x higher	> 3.0x higher
Compared to national average ($0.0001\mu\text{g}/\text{m}^3$)*	Below Average					Above Average				
BC	--	--	--	--	458,638 (10.4%)	23,444 (0.5%)	2,342,114 (53.2%)	294,565 (6.7%)	256,773 (5.8%)	1,024,523 (23.3%)
AB	--	--	--	--	2,307,115 (63.3%)	294,046 (8.1%)	713,020 (19.6%)	161,045 (4.4%)	63,509 (1.7%)	106,522 (2.9%)
SK	--	--	--	--	343,070 (33.2%)	13,368 (1.3%)	428,631 (41.5%)	50,249 (4.8%)	46,111 (4.5%)	151,952 (14.7%)
MB	--	--	--	--	839,055 (69.4%)	111,325 (9.2%)	157,430 (13.0%)	51,776 (4.3%)	19,565 (1.6%)	29,117 (2%)
ON	530,702 (4.1%)	116,580 (0.9%)	210,831 (1.6%)	70,415 (0.5%)	1,008,256 (7.8%)	3,368,353 (26.2%)	3,222,317 (25.1%)	845,401 (6.6%)	783,511 (6.1%)	2,695,455 (21.0%)
QC	--	--	--	--	1,050,012 (13.3%)	60,344 (0.8%)	1,461,606 (18.5%)	1,816,064 (23.0%)	1,123,101 (14.2%)	2,391,874 (30.3%)
NB	80,539 (10.7%)	11,310 (1.5%)	18,774 (2.5%)	13,346 (1.8%)	249,478 (33.2%)	13,492 (1.8%)	253,912 (33.8%)	21,983 (2.9%)	19,018 (2.5%)	69,319 (9.2%)
NS	--	--	--	--	274,048 (29.7%)	13,131 (1.4%)	438,968 (47.6%)	32,098 (3.5%)	36,117 (3.9%)	127,365 (13.8%)
PE	--	--	--	--	49,871 (3.6%)	2,464 (1.8%)	63,297 (45.1%)	4,809 (3.4%)	3,766 (2.7%)	15,997 (11.4%)
NL	--	--	--	--	202,329 (39.3%)	38,271 (7.4%)	168,695 (32.8%)	23,184 (4.5%)	29,580 (5.7%)	52,477 (10.2%)
NU	--	--	--	--	31,906 (100.0%)	--	--	--	--	--
NT	--	--	--	--	20,294 (48.9%)	576 (1.4%)	15,455 (37.3%)	2,570 (6.2%)	585 (1.4%)	1,982 (4.8%)
YT	--	--	--	--	6,991 (20.6%)	180 (0.5%)	17,463 (51.5%)	1,818 (5.4%)	1,577 (4.7%)	5,868 (17.3%)
CANADA	611,241 (1.8%)	127,890 (0.4%)	299,605 (0.7%)	83,761 (0.3%)	6,841,063 (20.4%)	3,938,994 (11.8%)	9,282,908 (27.7%)	3,305,562 (9.9%)	2,383,213 (7.1%)	6,672,451 (19.9%)

ASSOCIATED LIFETIME EXCESS CANCER RISK (per million people):
 RED = POTENTIAL LIFETIME EXCESS RISK IS GREATER THAN 1 PER MILLION PEOPLE

Health Canada CPF: No CPF	0.000033 to 0.00004	0.00004 to 0.00005	0.00005 to 0.000067	0.000067 to 0.0001	0.0001 to 0.00015	0.00015 to 0.0002	0.0002 to 0.00025	0.00025 to 0.0003	More than 0.0003	
California OEHHA CPF: 0.39	< 0.0003	0.0003 to < 0.0004	0.0004 to < 0.0005	0.0005 to < 0.0006	0.0006 to < 0.0009	0.0009 to < 0.0014	0.0014 to < 0.0018	0.0018 to < 0.0023	0.0023 to < 0.0027	> 0.0027
US EPA CPF: No CPF										

* measured at National Air Pollution Surveillance (NAPS) monitors in 2011
 CPF: Cancer Potency Factor